



NOVYRATE® EB PLUS

Scope	The increasing animal genetic potential puts a significant stress on the animals' digestive system, often leading to a suboptimal digestive process and nutrient absorption. The negative impact of sub-optimal feed utilization has a significant impact on the economical viability of the farm. Mono-butyric acid is an important dietary component that can help balance and stimulate a healthy intestinal microflora.
Description	Complementary feed. A synergetic combination of Esterified Butyrins, with immediate activity in the early part of the intestinal tract, and a prolonged effect towards the hindgut.
Components	Esterified butyric acid, flavour compounds, carrier and anti-caking.
Physical & Technical Specifications	<p>Physical appearance : Free flowing powder.</p> <p>Colour : white</p> <p>Odour : characteristic</p> <p>pH (10%) : 5,8 - 6,8</p> <p>Specific gravity : 0,66 - 0,76 kg/l</p> <p>Moisture (KF) : Max 3,5 %</p> <p>Colour change or variation does not affect performance.</p>
Application & Dosage	Piglets: 0.75 - 2 kg/T, Pigs (+ 25 kg): 0,25 - 0,75 kg/T, Sows: 1-2 kg/T, Broilers: 0,25 - 1 kg/T, Layers / Breeders: 0,5 - 0,75 kg/T, Turkeys: 0,5 - 1 kg/T
Packaging	Standard 25 kg export worthy bags on wooden fumigated pallets. Other net weights 1 kg*, 5 kg*, 10 kg, 20 kg, 750-1000kg, bulk on demand. *Marked unit packs are packed in an outer carton up to max. 25 kg.
Shelf Life	2 years when stored in a cool, well ventilated and dry environment out of direct sunlight in unopened packing.
Item Reference	10262

The information in this data sheet is to the best of our knowledge true and accurate. All instructions, recommendations and suggestions are made under reserve. Since the conditions of use are beyond our control, the manufacturer disclaims all liability for loss or damage suffered from the use of these data, suggestions or recommendations.